

HEEGAARD STRUCTURE RESPECTS COMPLICATED JSJ DECOMPOSITIONS

DAVID BACHMAN, RYAN DERBY-TALBOT, AND ERIC SEDGWICK

ABSTRACT. Let M be a 3-manifold with torus boundary components T_1 and T_2 . Let $\phi: T_1 \rightarrow T_2$ be a homeomorphism, M_ϕ the manifold obtained from M by gluing T_1 to T_2 via the map ϕ , and T the image of T_1 in M_ϕ . We show that if ϕ is “sufficiently complicated” then any incompressible or strongly irreducible surface in M_ϕ can be isotoped to be disjoint from T . It follows that every Heegaard splitting of a 3-manifold admitting a “sufficiently complicated” JSJ decomposition is an amalgamation of Heegaard splittings of the components of the JSJ decomposition.

1. INTRODUCTION

Let M be a (possibly disconnected) 3-manifold with homeomorphic boundary components F_1 and F_2 . Let $\phi: F_1 \rightarrow F_2$ be a homeomorphism, and let M_ϕ be the manifold obtained from M by gluing F_1 to F_2 via the map ϕ . Finally, let F denote the image of F_1 in M_ϕ . So-called incompressible and strongly irreducible surfaces have been seen to be vital to the study of 3-manifolds. A natural question is how such surfaces in M_ϕ are related to surfaces in M . In many cases it is known that with some assumption on the map ϕ , a class of such surfaces in M_ϕ can be isotoped to be disjoint from F . In other words, each surface in the class is isotopic into the original manifold M . When this conclusion follows, then we say F is a *barrier* for the class.

If F is separating and its genus is greater than one, then Lackenby [Lac04] and Souto [Sou] have independently shown that if ϕ is “sufficiently complicated” (in different contexts), then F is a barrier for all incompressible and strongly irreducible surfaces of bounded genus. Li [Li] also obtained a similar result, including the case that F is a (possibly non-separating) torus. In some cases it is even known that F is a barrier for all incompressible and strongly irreducible surfaces *regardless of their genus*. For example, this follows from [BSS06] when F is a separating torus in M_ϕ .

In this paper we look at the situation when the gluing surface is a torus T in M_ϕ that may be non-separating. Our main result is that the same conclusion holds: when ϕ is “sufficiently complicated” the surface T is a barrier for all incompressible and strongly irreducible surfaces, regardless of their genus. A consequence of this is that all Heegaard splittings of M_ϕ arise from Heegaard splittings of M in a natural way.

To prove our main result, the first challenge is to construct a reasonable definition of the term “sufficiently complicated” that depends only on the manifold M . This is accomplished in Section 2, where we construct a complexity for gluing maps between two torus boundary components of a 3-manifold called the *c-distance*, where c depends on the class of surfaces under consideration. We then establish our first theorem:

Theorem 1.1. *Let M be a compact, orientable, irreducible 3-manifold with incompressible boundary and let T_1 and T_2 be two torus components of ∂M . Let $\phi: T_1 \rightarrow T_2$ be a gluing map whose 1-distance is at least two. Then every closed, orientable, incompressible and strongly irreducible surface in M_ϕ can be isotoped to be disjoint from the image of T_1 in M_ϕ .*

That is, a sufficiently complicated gluing map between torus boundary components of a 3-manifold creates a barrier to all incompressible and strongly irreducible surfaces. The most interesting case of Theorem 1.1 is when M is connected, since similar results were previously obtained in [BSS06] for the case that M is disconnected. In the special case when $M \cong T^2 \times I$ this follows from [CS99].

The proof of Theorem 1.1 is given in Section 3, considering the incompressible and strongly irreducible cases separately. In Section 4 we review the definition of *generalized Heegaard splitting* and *amalgamation*, as presented in [Bac08]. We then extend the definition of *c-distance* to gluings along arbitrarily many tori, and put everything together to prove the following theorem about Heegaard splittings of certain 3-manifolds with non-trivial JSJ decompositions (see Section 4 for relevant definitions).

Theorem 1.2. *Let M be a closed, irreducible, orientable 3-manifold that admits a JSJ decomposition whose collection of JSJ tori has 1-distance at least 2. Then every Heegaard splitting of M is an amalgamation of Heegaard splittings of the components of the JSJ decomposition of M over the gluing maps.*

Theorem 1.2 greatly simplifies the Heegaard structure of 3-manifolds that are “sufficiently complicated” in the above sense. For example, despite being toroidal, such manifolds admit only finitely many Heegaard splittings of any given genus up to isotopy (see Corollary 4.12).

In the proof of Theorem 1.1 we show that ϕ need only to have 1-distance one to create a barrier to incompressible surfaces. In Section 5 we address the question of the existence of c -distance one gluing maps. There we show the following:

Theorem 1.3. *Let M be a compact, orientable, irreducible 3-manifold with incompressible boundary and let T_1 and T_2 be two torus components of ∂M . Let $\psi: T_1 \rightarrow T_2$ and $\sigma: T_2 \rightarrow T_2$ be homeomorphisms, where σ is Anosov. Then for each c there exists a positive integer N such that for $n \geq N$, the map $\sigma^n \psi$ has c -distance at least one.*

As above, the most interesting case is when M is connected since otherwise this result is implied by [Hat82]. Theorems 1.1 and 1.3 imply that the composition of any gluing map between two torus boundary components of some 3-manifold with a sufficiently high power of an Anosov map creates a barrier to all incompressible surfaces. We expect a similar result holds for strongly irreducible surfaces as well.

2. BOUNDARY SLOPES OF SURFACES THAT MEET THE 2-SKELETON NORMALLY

Let M be a compact, orientable, irreducible, triangulated 3-manifold consisting of one or two components¹. Presently we shall define what it means for a surface to meet the 2-skeleton of the triangulation normally. Such surfaces can then be broken up into *compatibility classes* with certain properties (see Definitions 2.5 and 2.7). Suppose T_1 and T_2 are torus components of ∂M . The main result of this section is an extension of Hatcher’s theorem [Hat82], [FO84] from slopes of essential surfaces to slopes of surfaces which meet the 2-skeleton of a triangulation normally. Namely, we show that each compatibility class $\mathcal{C}$ determines an element $\Phi_{\mathcal{C}}$ of $SL_2(\mathbb{Q})$ such that for each $S \in \mathcal{C}$, $\Phi_{\mathcal{C}}(\langle S \cap T_2 \rangle) = \langle S \cap T_1 \rangle$, where $\langle S \cap T_i \rangle$ is the slope of $S \cap T_i$. These maps will allow us to define the complexity for gluing maps $\phi: T_1 \rightarrow T_2$ referred to in Section 1.

¹The results for this section in the case that M has two components have been obtained previously in [Hat82] and [JS03]. While our focus is therefore on the case that M is connected, the results we establish here are general and include these previously obtained results in our own terminology, which we provide for completeness and ease of exposition in later sections.

Much of the terminology for this section is based on that of Jaco and Sedgwick [JS03]. Also, the reader is referred to [JT95] for a more elementary introduction to normal surface theory.

Fix a triangulation $\mathcal{T}$ of M that restricts to one-vertex triangulations on T_1 and T_2 . Such triangulations exist by [JS03].

Definition 2.1. A properly embedded arc in a 2-simplex of $\mathcal{T}$ is *normal* if it connects distinct 1-simplices. A loop on the boundary of a tetrahedron is normal if it consists of normal arcs. A *normal isotopy* of such a loop is an isotopy that restricts to an isotopy in each boundary simplex of the tetrahedron. It is well known that each normal loop consists of 3 or $4n$ normal arcs. The *complexity* of a normal loop which consists of 3 arcs is zero, and of a normal loop that consists of $4n$ arcs is $n - 1$.

Definition 2.2. A surface S in M is *normal with respect to the 2-skeleton* if it meets the boundary of every tetrahedron in a collection of normal loops. The *complexity* of such a surface is the maximum complexity of all such normal loops.

Examples 2.3.

- (1) Haken [Hak68] showed that surfaces that are both incompressible and ∂ -incompressible can be isotoped to be normal; i.e., to be complexity 0 surfaces, and in addition to meet each tetrahedron in disks.
- (2) It follows from work of Rubinstein [Rub95] and Stocking [Sto00] that Heegaard surfaces may be isotoped to be complexity at most 1, provided they are *strongly irreducible*. This condition means that any compressing disk on one side of such a Heegaard surface meets every compressing disk on the other, and that there is at least one such disk on each side.
- (3) It follows from work of the first author [Bac01], [Bac04], Coward [Cow], and Wilson [Wil08] that *∂ -strongly irreducible* surfaces can be isotoped to be complexity at most 1. Such surfaces have the property that any compressing or ∂ -compressing disk on one side meets every compressing and ∂ -compressing disk on the other.

Definition 2.4. Two surfaces F and G that are normal with respect to the 2-skeleton are *compatible* if on the boundary of each tetrahedron Δ , the normal loops $F \cap \partial\Delta$ and $G \cap \partial\Delta$ can be normally isotoped to be disjoint.




FIGURE 1. The six normal arc types in a 1-vertex triangulation of a torus.

It is well known that if F and G are compatible, and α and β are components of $F \cap \partial\Delta$ and $G \cap \partial\Delta$, respectively, of length larger than 3, then α and β are normally isotopic.

Definition 2.5. A maximal (with respect to inclusion) set of surfaces that are normal with respect to the 2-skeleton and pairwise compatible forms a *compatibility class*.

A choice of a normal loop of length larger than 3 on the boundary of each tetrahedron thus determines at most one compatibility class.

Definition 2.6. The *complexity* of a non-empty compatibility class is the maximum of the complexities of the surfaces contained in the class.

Following [JS03], a (not necessarily connected) normal curve in a one-vertex triangulation of a torus is parameterized by a free choice of normal coordinates $\{(x_1, x_2, x_3) \mid x_i \in \mathbb{N}\}$ in exactly one of the two triangles, forcing the normal coordinates in the other triangle to be equal, $x_1 = x_4, x_2 = x_5, x_3 = x_6$ (see Figure 1). Projectively, this set corresponds to the points with rational coordinates in the 2-simplex pictured in Figure 2.

A normal curve has *type* i if $x_i \leq x_j, j \neq i$. Of course, a normal curve with a non-unique minimal coordinate has more than one type, and will be all three types precisely when it consists entirely of trivial curves.

Definition 2.7. The *type* of a surface S in M is the pair of types of the boundary curves of S on T_1 and T_2 , respectively. A *typed compatibility class* is a compatibility class restricted to surfaces of the same type.

Remark 2.8. Since there are a bounded number of distinct normal isotopy classes of loops of bounded complexity on the boundary of a single tetrahedron, then it follows that there are a bounded number




FIGURE 2. The projective solution space has three types of normal curves.




FIGURE 3. Positive and negative intersections between normal arcs α and β .

of typed compatibility classes with complexity less than a specified bound. For instance, there are at most 3^{2t} typed compatibility classes of complexity zero surfaces, each determined by a pair of boundary types and a choice of a length 4 loop in each of the t tetrahedra.

Let α and β be normal arcs in an oriented triangle δ . Then α and β can be isotoped, keeping their boundaries fixed, so that they intersect in at most one point. We define the *normal sign* of the point $\alpha \cap \beta$, if it exists, as follows. There is at least one arc ϵ of the 1-skeleton of δ that connects α to β . Then we say $\alpha \cap \beta$ is

- *positive* if the orientation on ϵ induced by the orientation of δ points from β to α , and
- *negative* otherwise (see Figure 3).

Lemma 3.5 of [JS03] shows that normal isotopy and isotopy are equivalent for normal curves in a one-vertex triangulation of a torus. Moreover,




FIGURE 4. Type 3 curves can be oriented by orienting normal arcs as indicated.

Lemma 2.9. *Suppose that normal curves of the same type, α and β , have been normally isotoped to intersect minimally in a one-vertex triangulation of a torus. Then all intersection points between α and β have the same normal sign.*

Proof. Without loss of generality assume that α and β are both of type 3. Since they intersect minimally, each trivial component of α and β is disjoint from every other component. Let α' and β' be obtained by removing trivial components from α and β respectively. Then α' and β' do not possess normal arcs of the third type and can be consistently oriented by applying orientations to the remaining two arc types as in Figure 4.

Now consider an intersection between a normal arc of α' and a normal arc of β' . Since the curves are of type 3, the algebraic sign is equal to the normal sign of the intersection point (see Figure 5). We have already observed, however, that α' and β' intersect minimally, hence all intersections have the same algebraic sign. It follows that all intersection points have positive normal sign or all have negative normal sign.

□

Lemma 2.10. *Let A and B be surfaces meeting the 2-skeleton of M normally, in the same typed compatibility class, whose intersection with ∂M is contained in $T_1 \cup T_2$. Suppose A and B are normally isotoped to intersect minimally. Let $\alpha_i = A \cap T_i$ and $\beta_i = B \cap T_i$ designate their normal boundary curves, oriented by type (see Figure 4). Then the algebraic intersection numbers are related by $\#(\alpha_1, \beta_1) = -\#(\alpha_2, \beta_2)$.*

Proof. Consider a tetrahedron, Δ . Let α denote a component of $A \cap \partial\Delta$, and β a component of $B \cap \partial\Delta$. Since A and B intersect minimally, we may assume each normal arc of α and β is a straight line segment.




FIGURE 5. For intersections between oriented type 3 curves, the algebraic sign is equal to the normal sign.




FIGURE 6. Positive and negative intersections cancel as t increases through t_i .

As A and B are compatible, there is an isotopy α_t from α to a normal loop α' in $\partial\Delta$ that is disjoint from β . We can choose such an isotopy that interpolates the intersections with the 1-skeleton and so that for all t , each normal arc of α_t is a straight line segment. Furthermore, by adjusting the rate of interpolation, we may assume that the isotopy is in general position. Let $\{t_i\}$ denote the critical values of $|\alpha_t \cap \beta|$. It follows that for each i , $\alpha_{t_i} \cap \beta$ includes a point of the 1-skeleton. Let η_t denote the difference between the number of positive and negative intersection points of α_t and β , when their intersection is in general position.

Just before (or after) t_i , α_t meets β as in Figure 6. Here we see two intersections, one of each normal sign, of $\alpha_t \cap \beta$ which cancel as t increases through t_i . It follows that $\eta_{t_i - \epsilon} = \eta_{t_i + \epsilon}$. As $\alpha' \cap \beta = \emptyset$, we conclude η_t is zero for all non-critical t . It follows that there are an

equal number of positive and negative normal signs on the intersections of α and β on $\partial\Delta$.

Note that the induced orientations on an interior face δ are opposite from the two (not necessarily distinct) tetrahedra on either side of it. Therefore the normal sign of each intersection point is opposite with respect to each orientation. It follows that the sum of the normal intersections is zero when summed over only faces in the boundary. By Lemma 2.9, on each boundary component T_i , each intersection point between α_i and β_i has the same sign. Therefore the positive and negative intersections between A and B on $T_1 \cup T_2$ happen on different tori. Since they agree with the algebraic intersection numbers there, $\#(\alpha_1, \beta_1) = -\#(\alpha_2, \beta_2)$. $\square$

The above result is particularly relevant to the case that M is connected. In the case that M consists of two components (each having a torus boundary component), Lemma 2.10 implies that every surface in a given compatibility class (intersecting ∂M in $T_1 \cup T_2$ only) intersects each boundary torus in a unique slope.

Fix homology bases for T_1 and T_2 so that the orientation on T_1 agrees with the induced boundary orientation, and the orientation on T_2 disagrees with the induced boundary orientation. A *slope* is the isotopy class of a closed and connected essential curve in a torus. With our choice of basis for the homology of the torus, slopes are parameterized by $\widehat{\mathbb{Q}} = \mathbb{Q} \cup \{1/0\}$. If c is an embedded but disconnected curve that contains an essential curve, then we will define the *slope of c* , denoted $\langle c \rangle$, to be the isotopy class of one of its essential connected components. The *slope of a surface S* on a torus T is the slope of an essential component of its intersection with T . An element $\Phi \in SL_2(\mathbb{Q})$ acts on the set of slopes $\widehat{\mathbb{Q}}$ as a linear fractional transformation, i.e., $\Phi(q) = \frac{aq+b}{cq+d}$.

Lemma 2.11. *Let $\mathcal{C}$ be a typed compatibility class of surfaces that meet the 2-skeleton normally. Then there exists an element $\Phi_{\mathcal{C}} \in SL_2(\mathbb{Q})$ such that for all $S \in \mathcal{C}$ where $S \cap T_1 \neq \emptyset$, $S \cap T_2 \neq \emptyset$, and $S \cap \partial M \subset T_1 \cup T_2$, the element $\Phi_{\mathcal{C}}$ maps the slope of S on T_2 to the slope of S on T_1 , i.e. $\Phi_{\mathcal{C}}(\langle S \cap T_2 \rangle) = \langle S \cap T_1 \rangle$.*

Proof. There are two cases. In the first case, we assume that all surfaces $S \in \mathcal{C}$ meet T_2 in a single slope τ_2 . Since any pair of these surfaces can be isotoped to be disjoint on T_2 , it follows from Lemma 2.10 that they can also be isotoped to be disjoint on T_1 . Hence all surfaces $S \in \mathcal{C}$ have the same slope τ_1 on T_1 . With respect to the given bases on T_i choose a particular element $\Phi_{\mathcal{C}} \in SL_2(\mathbb{Q})$ so that $\Phi_{\mathcal{C}}(\tau_2) = \tau_1$.

In the second case, we suppose that there are two surfaces R and S in $\mathcal{C}$ which meet T_2 in distinct slopes, and hence in distinct slopes on T_1 . Let $\vec{r}_i$ and $\vec{s}_i$ be column vectors expressing the homology classes $[R \cap T_i]$ and $[S \cap T_i]$, respectively. Construct the matrices $\Psi_i = (\vec{r}_i \vec{s}_i)$. By Lemma 2.10 and because one of the chosen homology bases disagrees with the induced boundary orientations, $\det(\Psi_1) = \det(\Psi_2)$. Now let $\Phi_{\mathcal{C}} = \Psi_1 \Psi_2^{-1}$. Note that $\Phi_{\mathcal{C}}$ has determinant 1, and thus can be regarded as an element of $SL_2(\mathbb{Q})$. Restricting the equation $\Phi_{\mathcal{C}} \Psi_2 = \Psi_1$ to each column vector of Ψ_2 , we obtain $\Phi_{\mathcal{C}} \vec{r}_2 = \vec{r}_1$ and $\Phi_{\mathcal{C}} \vec{s}_2 = \vec{s}_1$, i.e., $\Phi_{\mathcal{C}}$ is a map between the homology classes of the boundaries of these surfaces.

We need to show this holds for any other surface $Q \in \mathcal{C}$. Suppose that Q meets the boundary with homology classes $\vec{q}_1$ and $\vec{q}_2$. Since $\vec{r}_1$ and $\vec{s}_1$ are independent, any other vector is uniquely determined by its intersections with them. We check that $\vec{q}_1$ and $\Phi_{\mathcal{C}} \vec{q}_2$ have the same number of intersections with $\vec{r}_1$:

$$\det(\Phi_{\mathcal{C}} \vec{q}_2, \vec{r}_1) = \det(\Phi_{\mathcal{C}} \vec{q}_2, \Phi_{\mathcal{C}} \vec{r}_2) = \det(\Phi_{\mathcal{C}}) \det(\vec{q}_2, \vec{r}_2) = \det(\vec{q}_1, \vec{r}_1),$$

by Lemma 2.10. The calculation for $\vec{s}_1$ is identical and thus $\Phi_{\mathcal{C}}$ maps $\vec{q}_2$ to $\vec{q}_1$, as desired.

Finally, in this second case, we regard $\Phi_{\mathcal{C}}$ as a linear fractional transformation so that it acts on slopes rather than homology classes. $\square$

We now define the notion of distance of a map that glues together two torus boundary components of a 3-manifold. To do this, we first define the Farey graph:

Definition 2.12. Let T be a torus. The *Farey graph* for T is the graph formed by taking a vertex for each slope on T , and connecting two vertices with an edge if simple closed curve representatives of the two corresponding slopes can be isotoped to intersect in one point on T . The distance between two slopes is taken to be the minimum length of a path in the Farey graph between the corresponding vertices.

Definition 2.13. Fix a number c . For each typed compatibility class $\mathcal{C}$ of surfaces of complexity at most c (of which there are a finite number by Remark 2.8), let $\Phi_{\mathcal{C}}$ be the element of $SL_2(\mathbb{Q})$ given by Lemma 2.11. Suppose $\phi: T_1 \rightarrow T_2$ is a gluing map. Then we say $\phi \Phi_{\mathcal{C}}$ has *distance* d if the minimum distance (in the Farey graph of T_2), over all slopes γ on T_2 , of γ and the slope of $\phi \Phi_{\mathcal{C}}(\gamma)$, is d . The *c-distance* of the gluing map ϕ is then the minimum, over all maps $\Phi_{\mathcal{C}}$, of the distances of $\phi \Phi_{\mathcal{C}}$.

Note that if the c -distance of ϕ is non-zero then each composition $\phi\Phi_C$ fixes no slope. Also, observe that if $c \leq c'$, then the c -distance of ϕ is greater than or equal to the c' -distance of ϕ .

3. PROOF OF THEOREM 1.1.

In this section we present the proof of Theorem 1.1. This is broken up into two cases, depending on whether the surfaces we are concerned with are incompressible or strongly irreducible.

3.1. The incompressible case.

Lemma 3.1. *Let X be a compact, irreducible, orientable 3-manifold with ∂X incompressible, if non-empty. Suppose S and T are closed, essential surfaces in X . Then S may be isotoped to be transverse to T with every component of $S \setminus N(T)$ incompressible in the respective submanifold of $X \setminus N(T)$.*

Proof. Isotope S so that it meets T transversally, and so that $|S \cap T|$ is minimal. Let α be a loop of $S \cap T$. Let D be a compressing disk for $S \setminus N(T)$. As S is incompressible in X , ∂D bounds a subdisk E of S , where necessarily $E \cap T \neq \emptyset$. As T is incompressible, every loop of $E \cap T$ must be inessential on T . We conclude $S \cap T$ contains some loop that is inessential on both surfaces. But then by a standard innermost disk argument we can lower $|S \cap T|$, a contradiction. Thus, $S \setminus N(T)$ must be incompressible in $X \setminus N(T)$.

We must now establish that each component of $S \setminus N(T)$ is incompressible. Choose the compressing disk C for a component of $S \setminus N(T)$ whose interior meets the other components in a minimal number of loops. Let α be such a loop that is innermost on C . Then α bounds a subdisk C' of C , and $C' \cap S = \partial C'$. As $S \setminus N(T)$ is incompressible, $\partial C'$ is inessential on $S \setminus N(T)$. But then by a similar innermost disk argument we may reduce the number of intersections of C with $S \setminus N(T)$, a contradiction. $\square$

Lemma 3.2. *Let M be a compact, orientable, irreducible 3-manifold. Let S be an orientable, connected, properly embedded, incompressible surface in M , such that each loop of ∂S is contained in a torus component of ∂M . Then either S is a boundary-parallel annulus or S is ∂ -incompressible.*

Proof. Suppose there is a ∂ -compression D' for S . Let T be the component of ∂M that meets D' . Then $D' \cap T$ is an arc, β . There are now two cases. Suppose first $\partial\beta$ connects two distinct loops of $S \cap T$. As T is a torus, these loops cobound an annulus A of T . Let A' denote the

disk obtained from A by removing a neighborhood of β . Then two parallel copies of D' , together with the disk A' , is a disk whose boundary is on S . As S is incompressible, we conclude S is a boundary-parallel annulus.

The second case is when $\partial\beta$ connects some loop of $S \cap T$ to itself. Then β , together with an arc of $S \cap T$, cobounds a subdisk A'' of T . The disk $D' \cup A''$ is then a compressing disk for S , a contradiction. $\square$

We now establish the incompressible case of Theorem 1.1. Recall that M is assumed to be a 3-manifold with incompressible boundary, $\phi: T_1 \rightarrow T_2$ is a gluing map between two torus components of ∂M , and M_ϕ is the resulting 3-manifold. Let T be the image of T_1 in M_ϕ . Fix a triangulation $\mathcal{T}$ of M as in Section 2. Let S be a closed, orientable, incompressible surface in M_ϕ .

By Lemma 3.1 we can isotope S so that $S' = S \setminus N(T)$ is incompressible in $M_\phi \setminus N(T) \cong M$. If, for some i , every component of S' that meets T_i can be isotoped into a neighborhood of T_i , then we can isotope S in M_ϕ to miss T entirely, and the result follows. Otherwise, by Lemma 3.2, for each i there is a component of S' that meets T_i and is both incompressible and ∂ -incompressible. Let S'' then be a (possibly disconnected) subsurface of S' , which is properly embedded in M , is incompressible, ∂ -incompressible, and meets both T_1 and T_2 .

By [Hak68] S'' can be isotoped to be a normal surface with respect to $\mathcal{T}$. In this position, S'' is a complexity zero surface in M such that $\langle \phi(S'' \cap T_1) \rangle = \langle S'' \cap T_2 \rangle$.

Let $\mathcal{C}$ denote the typed compatibility class of S'' . Then by Lemma 2.11, $\Phi_{\mathcal{C}}(\langle S'' \cap T_2 \rangle) = \langle S'' \cap T_1 \rangle$. It follows that $\phi\Phi_{\mathcal{C}}$ fixes the slope $S'' \cap T_1$, and thus the map ϕ has 0-distance (and hence 1-distance) zero. This concludes the proof of the incompressible case of Theorem 1.1.

3.2. The strongly irreducible case. When the surface S is strongly irreducible, we may appeal to the results of [BSS06]. Lemmas 3.3 and 3.5 below, taken from that paper, will play the roles that Lemmas 3.1 and 3.2 played in the previous subsection.

Lemma 3.3. [BSS06] *Let X be a compact, irreducible, orientable 3-manifold with ∂X incompressible, if non-empty. Suppose $X = V \cup_S W$, where S is a strongly irreducible Heegaard surface. Suppose further that X contains an incompressible, orientable, closed, non-boundary parallel surface T . Then either*

- *S may be isotoped to be transverse to T , with every component of $S \setminus N(T)$ incompressible in the respective submanifold of $X \setminus N(T)$, or*

- S may be isotoped to be transverse to T , with every component of $S \setminus N(T)$ incompressible in the respective submanifold of $X \setminus N(T)$ except for exactly one strongly irreducible component, or
- S may be isotoped to meet T in a single saddle tangency (and transverse elsewhere), with every component of $S \setminus N(T)$ incompressible in the respective submanifold of $X \setminus N(T)$.

When T is a torus we can rule out the third conclusion of this lemma, and thereby obtain the following corollary:

Corollary 3.4. *Let X be a compact, irreducible, orientable 3-manifold with ∂X incompressible, if non-empty. Suppose $X = V \cup_S W$, where S is a strongly irreducible Heegaard surface. Suppose further that X contains an essential torus T . Then S may be isotoped to be transverse to T with every component of $S \setminus N(T)$ incompressible in the respective submanifold of $X \setminus N(T)$ except for at most one strongly irreducible component.*

Proof. Suppose S has been isotoped to meet T in a single saddle tangency (and transverse elsewhere), with every component of $S \setminus N(T)$ incompressible in the respective submanifold of $X \setminus N(T)$. Then $\partial(S \setminus N(T))$ must be essential on S and both copies of T on $\partial N(T)$. But this means that the saddle intersection $S \cap T$ can be resolved in two different ways to obtain loops that are essential on the torus T , a contradiction. $\square$

Lemma 3.5. [BSS06] *Let M be a knot manifold. Let S be a separating, properly embedded, connected surface in M which is strongly irreducible, has non-empty boundary, and is not peripheral. Then either S is ∂ -strongly irreducible or ∂S is at most distance one from the boundary of some properly embedded surface which is both incompressible and ∂ -incompressible.*

In [BSS06] the term *knot manifold* was used to refer to a 3-manifold with a single boundary component, which is homeomorphic to a torus. However, the proof of Lemma 3.5 given in [BSS06] only uses the fact that each boundary component of the surface S is contained in a torus component of ∂M .

We now proceed with the proof of the strongly irreducible case of Theorem 1.1. As before, M is a 3-manifold with incompressible boundary, $\phi: T_1 \rightarrow T_2$ is a gluing map between two torus components of ∂M , M_ϕ is the resulting 3-manifold, and T is the image of T_1 in M_ϕ . Fix a triangulation $\mathcal{T}$ of M as in Section 2. Let S be a closed, strongly irreducible surface in M_ϕ .

By Corollary 3.4 we can isotope S so that every component of $S' = S \setminus N(T)$ is incompressible in $M_\phi \setminus N(T) \cong M$, with the exception of at most one strongly irreducible component. If, for some i , every component of S' that meets T_i can be isotoped into a neighborhood of T_i , then we can isotope S in M_ϕ to miss T entirely, and the result follows. Otherwise, either by Lemma 3.2 or by Lemma 3.5 there is a (possibly disconnected) surface S'' , which is properly embedded in M , is either incompressible and ∂ -incompressible or strongly irreducible and ∂ -strongly irreducible, and meets both T_1 and T_2 . Furthermore, the slope of $\phi(S'' \cap T_1)$ is at most distance one from the slope of $S'' \cap T_2$ in the Farey graph of T_2 .

By [Hak68] and [Bac01] the surface S'' can be isotoped to be a complexity zero or one surface in M . Let $\mathcal{C}$ denote the typed compatibility class of S'' . Then by Lemma 2.11, $\Phi_{\mathcal{C}}(\langle S'' \cap T_2 \rangle) = \langle S'' \cap T_1 \rangle$. It follows that $\phi\Phi_{\mathcal{C}}$ translates the slope of $S'' \cap T_2$ at most distance one in the Farey graph of T_2 , and thus the map ϕ has 1-distance zero or one. This concludes the proof of Theorem 1.1.

4. HEEGAARD SPLITTINGS OF 3-MANIFOLDS WITH SUFFICIENTLY COMPLICATED JSJ DECOMPOSITIONS ARE AMALGAMATIONS.

Generalized Heegaard splittings were first introduced by Scharlemann and Thompson in [ST94]. Here we review some of the basic definitions of this theory, as presented in [Bac08], and provide an application of Theorem 1.1.

Definition 4.1. A *compression body* is a 3-manifold which can be obtained by starting with some closed, orientable, connected surface, H , forming the product $H \times I$, attaching some number of 2-handles to $H \times \{0\}$, and capping off all resulting 2-sphere boundary components that are not contained in $H \times \{1\}$ with 3-balls. The boundary component $H \times \{1\}$ is referred to as ∂_+ . The rest of the boundary is referred to as ∂_- .

Definition 4.2. A *Heegaard splitting* of a 3-manifold M is an expression of M as a union $V \cup_H W$, where V and W are compression bodies that intersect in a transversally oriented surface $H = \partial_+ V = \partial_+ W$. If $V \cup_H W$ is a Heegaard splitting of M then we say H is a *Heegaard surface*.

Definition 4.3. A complex Σ is the *spine* of a Heegaard splitting $V \cup_H W$ of M if

- (1) Σ is the union of some subset of ∂M and a properly embedded graph in M .

- (2) H is parallel to the frontier of a neighborhood of Σ in M .

Definition 4.4. A *generalized Heegaard splitting (GHS)* $\mathcal{H}$ of a 3-manifold M is a pair of sets of pairwise disjoint, transversally oriented, connected surfaces, $\text{Thick}(\mathcal{H})$ and $\text{Thin}(\mathcal{H})$ (called the *thick levels* and *thin levels*, respectively), which satisfies the following conditions.

- (1) Each component M' of $M - \text{Thin}(\mathcal{H})$ meets a unique element H_+ of $\text{Thick}(\mathcal{H})$, and H_+ is a Heegaard surface in M' . Henceforth we will denote the closure of the component of $M - \text{Thin}(\mathcal{H})$ that contains an element $H_+ \in \text{Thick}(\mathcal{H})$ as $M(H_+)$.
- (2) As each Heegaard surface $H_+ \subset M(H_+)$ is transversally oriented, we can consistently talk about the points of $M(H_+)$ that are “above” H_+ or “below” H_+ . Suppose $H_- \in \text{Thin}(\mathcal{H})$. Let $M(H_+)$ and $M(H'_+)$ be the submanifolds on each side of H_- . Then H_- is below H_+ if and only if it is above H'_+ .
- (3) There is a partial ordering on the elements of $\text{Thin}(\mathcal{H})$ which satisfies the following: Suppose H_+ is an element of $\text{Thick}(\mathcal{H})$, H_- is a component of $\partial M(H_+)$ above H_+ , and H'_- is a component of $\partial M(H_+)$ below H_+ . Then $H_- > H'_-$.

Definition 4.5. A GHS $\mathcal{H}$ of M is *strongly irreducible* if every element $H_+ \in \text{Thick}(\mathcal{H})$ is strongly irreducible in $M(H_+)$.

Theorem 4.6. [Bac08], [ST94] *The thin levels of a strongly irreducible GHS are incompressible.*

Corollary 4.7. *Let M be a 3-manifold with incompressible boundary and let T_1 and T_2 be two torus components of ∂M . Let $\phi: T_1 \rightarrow T_2$ be a gluing map whose 1-distance is at least two, and let T be the image of T_1 in M_ϕ . Then T is isotopic to a thin level of every strongly irreducible GHS of M_ϕ .*

Proof. Let $\mathcal{H}$ be a strongly irreducible GHS of M_ϕ . By Theorem 4.6, the thin levels of $\mathcal{H}$ are incompressible. By Theorem 1.1, such surfaces can be isotoped to be disjoint from T . But then T lies in $M(H_+)$, for some thick level H_+ . As H_+ is strongly irreducible we may again apply Theorem 1.1 to conclude T can be isotoped to be disjoint from H_+ in $M(H_+)$. But then T lies in a compression body. The only incompressible surfaces in a compression body are isotopic to negative boundary components, and all such components are thin levels of $\mathcal{H}$. $\square$




FIGURE 7. The wider horizontal lines are the thick levels of a GHS, the narrower horizontal lines represent its thin levels, and the darker arcs represent the spine of its amalgamation.

In [Bac08] and [Lac08] it was shown how $\mathcal{H}$ determines a unique Heegaard splitting of M , called its *amalgamation*, generalizing a procedure of the same name defined in [Sch93]. The definition is illustrated schematically in Figure 7. The dark lines in the figure represent a complex which is the spine of a Heegaard splitting of the manifold M pictured. This complex has three parts. The first are those subsurfaces of ∂M that lie below thick levels. The second part is a collection of loops that lies entirely in compression bodies below thick levels. Finally, the third part is a collection of vertical arcs that connect the other two parts through compression bodies that lie above thick levels.

The following result about amalgamation is widely known; see e.g. [Bac08]:

Lemma 4.8. *Suppose M is a compact, orientable 3-manifold with incompressible boundary. Then every Heegaard splitting of M is the amalgamation of some strongly irreducible GHS.*

Definition 4.9. Suppose $\{M_i\}$ is a collection of 3-manifolds, and M is obtained from this collection by gluing pairs of boundary components by a collection of maps $\{\phi_j\}$. Let $\mathcal{H}$ be a GHS of M whose thin levels are precisely the images of the boundary components of $\{M_i\}$ in M . Then the thick levels of $\mathcal{H}$ are a set $\{H_i\}$, where H_i is a Heegaard surface in M_i . The amalgamation H of $\mathcal{H}$ is then said to be the *amalgamation of the Heegaard surfaces H_i over the maps $\{\phi_j\}$* .

Now we extend the notion of c -distance to gluings along multiple tori.

Definition 4.10. Let $\mathbf{T}$ denote a collection of pairwise disjoint, pairwise non-parallel, essential tori in a compact, orientable 3-manifold M . Let $\{T_i\}_{i=1}^n$ be an ordering of the elements of $\mathbf{T}$. Let M_0 denote the manifold obtained from M by cutting along all tori in $\mathbf{T}$. Let T'_i and T''_i denote the copies of T_i on ∂M_0 . Let M_i denote the manifold obtained from M_{i-1} by the gluing $\phi_i: T'_i \rightarrow T''_i$, defined so that the manifold $M_n \cong M$. Then we say the ordered collection $\{T_i\}$ has c -distance d if there is a triangulation of each manifold M_i so that the minimum of the c -distances of the maps ϕ_i is precisely d . The c -distance of $\mathbf{T}$ is the maximum c -distance taken over all of its orderings $\{T_i\}$.

Definition 4.11. Let M be a closed, irreducible, orientable 3-manifold. Then there exists a minimal, pairwise disjoint union $\mathbf{T}$ of tori embedded in M such that each component of $M - \mathbf{T}$ is Seifert fibered or atoroidal. This is called the *JSJ decomposition* of M , and $\mathbf{T}$ is called the *collection of JSJ tori* in M .

We are now ready to prove Theorem 1.2.

Proof. Let H be a Heegaard surface of M , and let $\mathbf{T}$ denote the collection of JSJ tori in M . Let $\{T_i\}_{i=1}^n$ be an ordering of the JSJ tori realizing the 1-distance of $\mathbf{T}$, and let $\{M_i\}$ be the corresponding submanifolds as given in Definition 4.10. Now for some i , H is an amalgamation of splittings of the components of the manifold M_i , where at worst $i = n$. Let m be the smallest value of i such that H arises from an amalgamation of splittings of the components of M_m over the maps $\{\phi_i\}_{i=m+1}^n$. We claim $m = 0$, and thus the result follows.

Let H_m denote the (possibly disconnected) Heegaard surface of M_m whose amalgamation over the maps $\{\phi_i\}_{i=m+1}^n$ is H . By Lemma 4.8 H_m is the amalgamation of a strongly irreducible GHS $\mathcal{H}_m$ of M_m .

By definition, M_m is obtained from M_{m-1} by gluing two components of ∂M_{m-1} by a map $\phi_m : T'_m \rightarrow T''_m$ whose 1-distance is at least two. The image of T'_m in M_m is an essential torus T_m . By Corollary 4.7, T_m is a thin level of $\mathcal{H}_m$. We may thus obtain a strongly irreducible GHS $\mathcal{H}_{m-1}$ of M_{m-1} from $\mathcal{H}_m$ by replacing T_m in $\text{Thin}(\mathcal{H}_m)$ by the set $\{T'_m, T''_m\}$.

Let H_{m-1} denote the amalgamation of $\mathcal{H}_{m-1}$. Then by definition, H_m is the amalgamation of the components of H_{m-1} along the map ϕ_m . It follows that H is the amalgamation of the components of H_{m-1} along the maps $\{\phi_i\}_{i=m}^n$. We have thus contradicted the minimality of our choice of m . $\square$

A consequence of Theorem 1.2 is that 3-manifolds admitting “sufficiently complicated” JSJ decompositions in the above sense have only finitely many Heegaard splittings of a given genus, and thus do not satisfy the converse of the generalized Waldhausen Conjecture.

Corollary 4.12. *Let M be a closed, irreducible, orientable 3-manifold that admits a JSJ decomposition whose collection of JSJ tori has 1-distance at least 2. Then for every positive integer g , M admits at most finitely many Heegaard splittings of genus g up to isotopy.*

Proof. By Theorem 1.2, every Heegaard splitting of M is an amalgamation of Heegaard surfaces of the components of the JSJ decomposition of M . Since the amalgamation of Heegaard surfaces is unique up to isotopy, we only need to verify that each component of the JSJ decomposition admits finitely many Heegaard surfaces of a given genus up to isotopy. This follows from [LM91] and [Sch95] in the case that a component is Seifert fibered, and from [Joh95] in the case a component is atoroidal. $\square$

5. GLUING MAPS WITH c -DISTANCE AT LEAST ONE

In this final section, we prove Theorem 1.3. Recall that M is a 3-manifold with tori T_1 and T_2 contained in its boundary. To prove the theorem, we exploit the fact that homeomorphisms between T_1 and T_2 can be considered as matrices in $SL_2(\mathbb{Z})$. Therefore, to show that a gluing map $\sigma^n \psi$ has c -distance at least one, it suffices to show that for each c , the corresponding element in $SL_2(\mathbb{Z})$ composed with each Φ_c from Lemma 2.11 fixes no slopes. We do this by first establishing several linear algebraic results.

Definition 5.1. Let $L \in SL_2(\mathbb{Q})$, written in matrix form. Then the *denominator* of L , denoted $d(L)$, is the least integer $d \geq 1$ so that all entries of the matrix dL are integers.

Definition 5.2. Let $L \in SL_2(\mathbb{R})$, and let $\vec{v} = \begin{bmatrix} v_0 \\ v_1 \end{bmatrix}$ be an eigenvector of L . Then $r = v_0/v_1 \in \widehat{\mathbb{R}} = \mathbb{R} \cup \{\infty\}$ is an *eigenslope* of L . An eigenslope r is *rational* if $r \in \widehat{\mathbb{Q}} = \mathbb{Q} \cup \{\infty\}$.

Lemma 5.3. Suppose that $L \in SL_2(\mathbb{Q})$, r and s are non-zero integers, and $\vec{u}$ and $\vec{v}$ are vectors with coprime integer entries so that $Lr\vec{u} = s\vec{v}$. Then $d(L) \geq |s/r| \geq 1/d(L)$, where $d(L)$ is the denominator of L .

Proof. Suppose that $Lr\vec{u} = s\vec{v}$, where $\vec{u}$ and $\vec{v}$ have coprime integer entries. We first show that $|s/r| \geq 1/d(L)$. Notice $Lr\vec{u} = r/d(L)(d(L)L\vec{u}) = s\vec{v}$, where $d(L)L\vec{u}$ must have integer entries. Since $\vec{v}$ has coprime entries, $r/d(L)$ must be a divisor of s . It follows that $|s| \geq |r|/d(L)$, and hence $|s/r| \geq 1/d(L)$.

Now we show that $d(L) \geq |s/r|$. We have that $r\vec{u} = L^{-1}s\vec{v}$. Note that $d(L^{-1}) = d(L)$. It follows from above that $|r/s| \geq 1/d(L)$, and hence $d(L) \geq |s/r|$ as desired. $\square$

Lemma 5.4. Let $L \in SL_2(\mathbb{Q})$ so that $|\text{trace}(L)| < 2/d(L)$ or $|\text{trace}(L)| > 2d(L)$. Then L has no rational eigenslopes.

Proof. We prove the contrapositive. Suppose that L has a rational eigenslope written r/s , where r and s are coprime integers. Then $\vec{u} = \begin{bmatrix} r \\ s \end{bmatrix}$ is an eigenvector for L . Hence $L\vec{u} = q\vec{u}$ where q is the eigenvalue corresponding to $\vec{u}$. Since $L \in SL_2(\mathbb{Q})$ it follows that $q \in \mathbb{Q}$. By Lemma 5.3, we have $d(L) \geq |q| \geq 1/d(L)$. The eigenvalues of L are q and $1/q$, either both positive or both negative. In particular, $|q + 1/q| = |q| + |1/q|$. Moreover, $d(L) \geq |q| \geq 1/d(L)$ implies that $1/d(L) \leq |1/q| \leq d(L)$. Hence $2d(L) \geq |q| + |1/q| = |q + 1/q| = |\text{trace}(L)| \geq 2/d(L)$. $\square$

Lemma 5.5. Let $J \in SL_2(\mathbb{Z})$ and $K \in SL_2(\mathbb{Q})$ so that $\text{trace}(J) > 2$. Then there exists an $N \in \mathbb{N}$ so that for all $n \geq N$, either $|\text{trace}(J^n K)| < 2/d(J^n K)$ or $|\text{trace}(J^n K)| > 2d(J^n K)$. In particular, $J^n K$ has no rational eigenslopes.

Proof. Let t_n be the trace of $J^n K$. Since $|\text{trace}(J)| > 2$, J is conjugate to a matrix W in $SL_2(\mathbb{R})$ of the form:

$$W = \begin{bmatrix} w & 0 \\ 0 & \frac{1}{w} \end{bmatrix}.$$

Without loss of generality, assume $w > 1$ (the cases that $w < 1$ or w is negative are similar). Assume that C is the conjugation matrix for J , i.e. $C^{-1}JC = W$. Now set $Z = C^{-1}KC$. Thus Z is a fixed matrix in $SL_2(\mathbb{R})$:

$$Z = \begin{bmatrix} z_1 & z_2 \\ z_3 & z_4 \end{bmatrix}.$$

Since trace is invariant under conjugation, we have

$$\begin{aligned} t_n &= \text{trace}(J^n K) = \text{trace}(C^{-1}J^n C \cdot C^{-1}KC) \\ &= \text{trace}(C^{-1}JC \cdots C^{-1}JC \cdot C^{-1}KC) = \text{trace}(W^n Z). \end{aligned}$$

Now, since

$$W^n = \begin{bmatrix} w^n & 0 \\ 0 & \frac{1}{w^n} \end{bmatrix},$$

it follows that

$$\begin{aligned} t_n &= \text{trace}(W^n Z) = \text{trace} \left(\begin{bmatrix} w^n & 0 \\ 0 & \frac{1}{w^n} \end{bmatrix} \begin{bmatrix} z_1 & z_2 \\ z_3 & z_4 \end{bmatrix} \right) \\ &= \text{trace} \left(\begin{bmatrix} z_1 w^n & z_2 w^n \\ \frac{z_3}{w^n} & \frac{z_4}{w^n} \end{bmatrix} \right) = z_1 w^n + \frac{z_4}{w^n}. \end{aligned}$$

Let k be a positive constant. First, assume that $z_1 = 0$. If $z_4 = 0$ as well, then $t_n = 0$ for any n ; otherwise a high enough power of N gives that for $n \geq N$, $|t_n| = \left| \frac{z_4}{w^n} \right| < 2k$. In either case, the trace is certainly less than $2k$.

Now assume that $z_1 \neq 0$. Then clearly, for large enough n , $|t_n| = \left| z_1 w^n + \frac{z_4}{w^n} \right|$ can be taken to be greater than $2k$. The lemma follows by letting $k = 1/d(J^n K)$ in the former case and $k = d(J^n K)$ in the latter, and finally applying Lemma 5.4.

□

We now prove Theorem 1.3.

Proof. Let M be a 3-manifold and let T_1 and T_2 be two torus components of ∂M . Let $\psi: T_1 \rightarrow T_2$ and $\sigma: T_2 \rightarrow T_2$ be homeomorphisms. We can use these maps to glue M along T_1 and T_2 and obtain the manifold $M_{\sigma^n \psi}$. Assume that σ is Anosov.

For each $i = 1, 2$, fix a homology basis for T_i as in Section 2. As homeomorphisms induce maps between homology, for convenience we can identify the homeomorphisms ψ and σ with their induced maps so that $\psi: H_1(T_1) \rightarrow H_1(T_2)$ and $\sigma: H_1(T_2) \rightarrow H_1(T_2)$ are regarded as elements of $SL_2(\mathbb{Z})$, where σ is hyperbolic (in particular, $|\text{trace}(\sigma)| > 2$).

Fix a positive integer c , and let $\mathcal{C}$ be a typed compatibility class of complexity at most c . Let $\Phi_{\mathcal{C}}$ be the element in $SL_2(\mathbb{Q})$ given by Lemma 2.11. The element $\sigma^n \psi \Phi_{\mathcal{C}}$ is thus in $SL_2(\mathbb{Q})$, and fixes a slope on T_2 if and only if $\sigma^n \psi \Phi_{\mathcal{C}}$ has an eigenvector $\vec{v}$ in $H_1(T_2)$. In other words, $\sigma^n \psi \Phi_{\mathcal{C}}$ has distance zero if and only if $\sigma^n \psi \Phi_{\mathcal{C}}$ has a rational eigenslope.

By taking $J = \sigma$ and $K = \psi \Phi_{\mathcal{C}}$ and applying Lemma 5.5, we obtain an integer $N_{\mathcal{C}}$ such that for all $n \geq N_{\mathcal{C}}$, $\sigma^n \psi \Phi_{\mathcal{C}}$ has no rational eigenslopes. Hence, $\sigma^n \psi \Phi_{\mathcal{C}}$ cannot have distance zero. Following Remark 2.8, there are a finite number of typed compatibility classes of complexity at most c . By taking N to be the maximum over all integers $N_{\mathcal{C}}$ obtained as above, we have that $n \geq N$ implies that $\sigma^n \psi$ cannot have c -distance zero.

□

REFERENCES

- [Bac01] David Bachman. Heegaard splittings with boundary and almost normal surfaces. *Topology Appl.*, 116:153–184, 2001.
- [Bac04] David Bachman. Erratum to: “Heegaard splittings with boundary and almost normal surfaces” [Topology Appl. **116** (2001), no. 2, 153–184]. *Topology Appl.*, 143(1-3):279, 2004.
- [Bac08] David Bachman. Connected Sums of Unstabilized Heegaard Splittings are Unstabilized. *Geometry & Topology*, 12:2327–2378, 2008.
- [BSS06] David Bachman, Saul Schleimer, and Eric Sedgwick. Sweepouts of amalgamated 3-manifolds. *Algebr. Geom. Topol.*, 6:171–194, 2006.
- [Cow] Alexander Coward. Algorithmically detecting the bridge number of hyperbolic knots. Preprint. [arXiv:0710.1262](https://arxiv.org/abs/0710.1262).
- [CS99] Daryl Cooper and Martin Scharlemann. The structure of a solvmanifold’s Heegaard splittings. In *Proceedings of 6th Gökova Geometry-Topology Conference*, volume 23, pages 1–18, 1999.
- [FO84] W. Floyd and U. Oertel. Incompressible surfaces via branched surfaces. *Topology*, 23(1):117–125, 1984.
- [Hak68] Wolfgang Haken. Some results on surfaces in 3-manifolds. In *Studies in Modern Topology*, pages 39–98. Math. Assoc. Amer. (distributed by Prentice-Hall, Englewood Cliffs, N.J.), 1968.
- [Hat82] A. E. Hatcher. On the boundary curves of incompressible surfaces. *Pacific J. Math.*, 99(2):373–377, 1982.
- [Joh95] Klaus Johannson. *Topology and combinatorics of 3-manifolds*. Number 1599 in Lecture Notes in Mathematics. Springer, 1995.

- [JS03] William Jaco and Eric Sedgwick. Decision problems in the space of Dehn fillings. *Topology*, 42(4):845–906, 2003.
- [JT95] William Jaco and Jeffrey L. Tollefson. Algorithms for the complete decomposition of a closed 3-manifold. *Illinois J. Math.*, 39(3):358–406, 1995.
- [Lac04] Marc Lackenby. The Heegaard genus of amalgamated 3-manifolds. *Geom. Dedicata*, 109:139–145, 2004.
- [Lac08] Marc Lackenby. An algorithm to determine the Heegaard genus of simple 3-manifolds with nonempty boundary. *Algebr. Geom. Topol.*, 8(2):911–934, 2008.
- [Li] Tao Li. On the Heegaard splittings of amalgamated 3-manifolds. Preprint. [arXiv:0701395](https://arxiv.org/abs/0701395).
- [LM91] Martin Lustig and Yo'av Moriah. Nielsen equivalence in Fuchsian groups and Seifert fibered spaces. *Topology*, 30(2):191–204, 1991.
- [Rub95] J. Hyam Rubinstein. Polyhedral minimal surfaces, Heegaard splittings and decision problems for 3-dimensional manifolds. In *Proceedings of the Georgia Topology Conference*, pages 1–20, 1995.
- [Sch93] Jennifer Schultens. The classification of Heegaard splittings for (compact orientable surface) $\times S^1$. *Proc. London Math. Soc. (3)*, 67(2):425–448, 1993.
- [Sch95] Jennifer Schultens. Heegaard splittings of Seifert fibered spaces with boundary. *Trans. Amer. Math. Soc.*, 347(7):2533–2552, 1995.
- [Sou] Juan Souto. Distances in the curve complex and Heegaard genus. Preprint. Available at www.picard.ups-tlse.fr/%7Esouto/Heeg-genus.pdf.
- [ST94] Martin Scharlemann and Abigail Thompson. Thin position for 3-manifolds. In *Geometric topology (Haifa, 1992)*, volume 164 of *Contemp. Math.*, pages 231–238. Amer. Math. Soc., Providence, RI, 1994.
- [Sto00] Michelle Stocking. Almost normal surfaces in 3-manifolds. *Transactions of the American Mathematical Society*, 352:171–207, 2000.
- [Wil08] Robin Wilson. Meridional almost normal surfaces in knot complements. *Algebraic & Geometric Topology*, 8:1717–1740, 2008.

PITZER COLLEGE

E-mail address: bachman@pitzer.edu

QUEST UNIVERSITY CANADA

E-mail address: rdt@questu.ca

DEPAUL UNIVERSITY

E-mail address: esedgwick@cdm.depaul.edu